

Género y especie	WH 44	WH 45	WH 46	WH 55	WH 56	WH 58	WH 65	XII 16
Subfamilia Catocalinae								
<i>Catocala conjuncta</i> (Esper, 1786)	.	.	.	+	.	.	.	.
<i>Catocala conversa</i> (Esper, 1787)	.	+	.	+	x	.	.	.
<i>Catocala dilecta</i> (Hübner, 1808)	.	.	+	.	.	.	.	.
<i>Catocala elocata</i> (Esper, 1786)	.	.	.	+	+	+	.	+
<i>Catocala nymphaea</i> (Esper, 1788)	+	.	.	.	+	.	.	.
<i>Catocala nymphagoga</i> (Esper, 1787)	.	.	.	+	+	.	.	.
<i>Catocala optata</i> (Godart, 1824)	.	.	.	+	+	.	.	.
<i>Catocala promissa</i> (D. & Schif., 1775)	.	.	.	.	+	.	.	.
<i>Catocala sponso</i> (Hübner, 1808)	.	+	.	.	.	.	.	.
<i>Clytie sancta</i> (Staud., 1897)	.	.	.	.	.	.	.	+
<i>Dysgonia algira</i> (L., 1767)	.	.	.	+	+	.	.	.
<i>Dysgonia torrida</i> (Guenée, 1852)	.	.	.	.	.	.	.	+
<i>Euclidia glyphica</i> (L., 1758)	.	.	.	+	+	.	.	.
<i>Minucia lunaris</i> (D. & Schif., 1775)	.	+	□	+	+	.	+	.
<i>Ophiura tirhaca</i> (Cramer, 1780)	+	.	.	.	.	.	.	.
<i>Synthymia fixa</i> (F., 1775)	.	.	.	.	+	.	.	+
n.º de spp. por cuadrícula	2	3	2	8	10	1	1	
Número total de especies: 44.								
Subfamilia Cuculliinae								
<i>Agrochola helvola</i> (L., 1758)	.	.	.	.	+	.	.	.
<i>Agrochola lychnidis</i> (D. & Schif., 1775)	.	.	.	.	+	.	.	.
<i>Agrochola inercialis</i> (Staud., 1873)	.	.	.	.	+	.	.	.
<i>Allophyes alfaroi</i> Agenjo, 1952	.	.	.	.	+	.	.	.
<i>Amephana aurita</i> (F., 1787)	.	.	.	.	.	.	.	+
<i>Ammopolia wizenmanni</i> (Standfuss, 1850)	.	.	.	+	+	.	.	.
<i>Aporophyla nigra</i> (Haworth, 1810)	.	.	.	.	+	.	.	.
<i>Catophasia almoravida</i> (Graslin, 1863)	+	+	.	.	.	.	.	.
<i>Cleonymia baetica</i> (Rambur, 1839)	.	.	.	+	.	.	.	+
<i>Cleonymia yvanii</i> (Duponchel, 1833)	.	+	.	.	.	.	.	.
<i>Conistra ligula</i> (Esper, 1791)	+	.	.	.	.	.	.	.
<i>Conistra vacini</i> (L., 1761)	.	+	.	.	+	.	.	.
<i>Cucullia caninae</i> (Rambur, 1833)	.	.	.	.	+	.	.	.
<i>Cucullia santolinae</i> (Rambur, 1831)	+	+	.	.	.	.	.	.
<i>Cucullia verbasci</i> (L., 1758)	.	.	.	.	+	.	.	.
<i>Dryobota labecula</i> (Esper, 1788)	.	.	.	+	.	.	.	.
<i>Dryobotodes cerris</i> (Boisduval, 1840)	.	.	.	.	+	.	.	.
<i>Gripesia aprilinea</i> (L., 1758)	.	.	.	+	.	.	.	.
<i>Jodia croceago</i> (D. & Schif., 1775)	.	.	.	.	+	.	+	.
<i>Leucochlaena oditis</i> (Hübner, 1832)	.	.	.	.	+	.	.	.
<i>Litophane semibrunnea</i> (Haworth, 1809)	.	+	.	.	.	.	.	.
<i>Lophoterges millierei</i> (Staud., 1870)	.	.	□	.	.	.	.	.
<i>Metopoceras felicina</i> (Donzel, 1844)	.	.	.	.	.	.	.	+
<i>Metopoceras khalildja</i> Oberthür, 1884	.	.	.	+	.	.	.	.
<i>Omphalophana serrata</i> (Treitschke, 1826)	.	.	.	+	+	.	.	.
<i>Omphaloscelis lunosa</i> (Haworth, 1809)	.	.	.	+	+	.	.	.
<i>Polymixis argillacengo</i> (Hübner, 1822)	.	.	.	+	+	.	.	.